

Literature 10

- Alacs, E. A., Georges, A., FitzSimmons, N. N., & Robertson, J. 2009. DNA detective: a review of molecular approaches to wildlife forensics. *Forensic Science, Medicine and Pathology*. <https://doi.org/10.1007/s12024-009-9131-7>
- Alibardi, L. 2012. Immunolocalization of keratin-associated beta-proteins (beta-keratins) in the regenerating lizard epidermis indicates a new process for the differentiation of the epidermis in lepidosaurians. *Journal of Morphology* **273**(11): 1272–1279. <https://doi.org/10.1002/jmor.20057>
- Alibardi, L. & Rogers, G. 2015. Observations on fur development in *Echidna* (Monotremata, Mammalia) indicate that spines precede hairs in ontogeny. *The Anatomical Record* **298**: 761–770. <https://doi.org/10.1002/ar.23081>
- Allen, J. A., Lang, H. & Chapin, J. P. 1917. The American Museum Congo Expedition collection of bats. *Bulletin of the American Museum of Natural History* **37**: 405–563. <https://doi.org/10.5962/bhl.title.82385>
- Amerasinghe, F. P. 1983. The structure and identification of the hairs of the mammals of Sri Lanka. *Ceylon Journal of Science (Biological Science)* **16**(1–2): 76–125.
- Amman, B. R., Owen, R. D. & Bradley, R. D. 2002. Utility of hair structure for taxonomic discrimination in bats, with an example from the bats of Colorado. *Occasional Papers, Museum of Texas Technical University* **216**: 1–14.
- Ancillotto, L. & Mori, E. 2017. Adaptive significance of coat colouration and patterns of Sciuromorpha (Rodentia). *Ethology, Ecology & Evolution* **29**(3): 214–254. <https://doi.org/10.1080/03949370.2016.1145146>
- Appleyard, H. M. 1960. *Guide to the identification of animal fibres*. Wool Industries Research Association, Leeds, England, 188 pp.
- Aristotle. BC 300. *Parts of Animals, Movement of Animals, Progression of Animals (original: De Partibus Animalium)*. Translated by A. L. Peck & E. S. Forster. Loeb Classical Library 323, Cambridge, MA, Harvard University Press, 580 pp.
- Arita, H. & Aranda, M. 1987. *Técnicas para el estudio y clasificación de los pelos*. Instituto Nacional de Investigaciones sobre Recursos Bióticos, Xalapa, Veracruz. 21 pp.
- Armitage, K. B. 2009. Fur color diversity in marmots. *Ethology, Ecology & Evolution* **21**: 183–194. <https://doi.org/10.1080/08927014.2009.9522474>
- Arslan, A., Kryštufek, B., Matur, F. & Zima, J. 2016. Review of chromosome races in blind mole rats (*Spalax* and *Nannospalax*). *Folia Zoologica* **65**(4): 249–301.
- Asher, R. J. & Helgen, K. M. 2010. Nomenclature and placental mammal phylogeny. *BMC Evolutionary Biology* **10**: 102. <https://doi.org/10.1186/1471-2148-10-102>
- Bahuguna, A. 2011. *Trichotaxonomy of species of the families Sciuridae, Viverridae, Mustelidae, Herpestidae and Tragulidae*. Kolkata, Zoological Survey of India, 297 pp.
- Baker, B. W. 1980. Hair-catchers aid in identifying mammalian predators of ground-nesting birds. *Wildlife Society Bulletin* **8** (3): 257–259.
- Ballesteros-Duperón, E., Virgós, E., Moleón, M., Barea-Azcón, J. & Gil-Sánchez, J. M. 2014. How accurate are coat traits for discriminating wild and hybrid forms of *Felis silvestris*? Mammalia. De Gruyter 1–10. <https://doi.org/10.1515/mammalia-2013-0026>
- Banks, W. J. 1981. *Applied veterinary histology*. William and Wilkins, Baltimore, London, 292–346.

- Beja-Pereira, A., Oliviera, R., Alves, P. C., Schwartz, M. K. & Luikart, G. 2009. Advancing ecological understandings through technological transformations in noninvasive genetics. *Molecular Ecology Resources* **9**: 1279–1301. <https://doi.org/10.1111/j.1755-0998.2009.02699.x>
- Ben-David, M. & Flaherty, E. 2012. Stable isotopes in mammalian research: a beginner's guide. *Journal of Mammalogy* **93**(2): 312–328. <https://doi.org/10.1644/11-MAMM-S-166.1>
- Benedict, F. A. 1957. Hair structure as a generic character in bats. *University of California Publications, Zoology* **59**: 285–548.
- Benton, M. J. 1991. *The rise of the mammals*. Crescent Books, New York, pp. 24–45.
- Berger, L. R., Pickering, R., Kuhn, B., Backwell, L. R., Hancox, P. J., Kramers, J. D. & Boshoff, P. A. 2009. A Mid-Pleistocene in situ fossil brown hyaena (*Parahyaena brunnea*) latrine from Gladysvale Cave, South Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* **279**: 131–136. <https://doi.org/10.1016/j.palaeo.2009.05.004>
- Bertazzo, A., Biasiolo, M., Costa, C. V. L., de Stefani, E. C. & Allegrì, G. 2000. Tryptophan in human hair: correlation with pigmentation. *Il Farmaco* **55**: 521–525. [https://doi.org/10.1016/S0014-827X\(00\)00038-0](https://doi.org/10.1016/S0014-827X(00)00038-0)
- Bertolino, S., Cordero di Montezemolo, N., Preatoni, D. G., Wauters, L. A. & Martinoli, A. 2014. A grey future for Europe: *Sciurus carolinensis* is replacing native red squirrels in Italy. *Biological Invasions* **16**: 53–62. <https://doi.org/10.1007/s10530-013-0502-3>
- Bihari, Z., Csorba, G. & Heltai, M. (eds) 2007. *Magyarország emlőseinek atlasza*. In: Kosuth Természettár, Kossuth Kiadó, Budapest, 360 pp. [In Hungarian]
- Bisbing, R. E. 1982. *The forensic identification and association of human hair*. In: Saferstein, R. (ed.): Forensic science handbook. Prentice Hall, Englewood Cliffs, New Jersey, pp. 184–221.
- Blazej, A., Galatík, A., Galatík, J., Krul, Z. & Mládek, M. 1989. *Atlas of microscopic structures of fur skins 1*. Elsevier Science Publishers B. V., New York, Amsterdam.
- Bonnichsen, R., Hodges, L., Rema, W., Field, K. G., Kirner, D. L. & Taylor, R. E. 2001. Methods for the study of ancient hair: radiocarbon dates and gene from individual hairs. *Journal of Archaeological Science* **28** (7): 775–785. <https://doi.org/10.1006/jasc.2000.0624>
- Bragulla, H. H. & Homberger, D. G. 2009. Structure and functions of keratin proteins in simple, stratified, keratinized and cornified epithelia. *Journal of Anatomy* **214**: 516–559. <https://doi.org/10.1111/j.1469-7580.2009.01066.x>
- Brunner, H. & Coman, B. J. 1974. *The identification of Mammalian hair*. Inkatha Press, Melbourne.
- Brunner, H. & Triggs, B. 2002. *Hair ID: an interactive tool for identifying Australian mammalian hair*. CSIRO Publishing, Collingwood.
- Bullen, R. D. & McKenzie, N. L. 2008. Aerodynamic cleanliness in bats. *Australian Journal of Zoology* **56**: 281–296. <https://doi.org/10.1071/ZO08059>
- Caro, T. 2009. Contrasting coloration in terrestrial mammals. *Philosophical transactions of the Royal Society B* **364**: 537–548. <https://doi.org/10.1098/rstb.2008.0221>
- Chakraborty, R., Chakraborty, S. & De, J. K. 1996. Identification of dorsal guard hairs of five Indian species of the genus *Panthera* Oken (Carnivora: Felidae). *Mammalia* **60**: 473–480. <https://doi.org/10.1515/mamm-1996-0312>
- Chakraborty, R. & De, J. K. 2010. *Atlas on hairs of Indian Mammals, Part I: Carnivora*. Published by the Director, Zoological Survey India, Kolkata, 144 pp.

- Chehébar, Cl. & Martín, S. 1989. Guía para el reconocimiento microscópico de los pelos mamíferos de la Patagonia. *Donana, Acta Vertebrata* **16**(2): 247–291.
- Chernova, O. F. 2002. Architectonic and diagnostic significance of hair cuticle. *Biology Bulletin* **29**: 238–247. <https://doi.org/10.1023/A:1015482430438>
- Chernova, O. F. 2003. Architectonic and diagnostic significance of hair cortex and medulla. *Biology Bulletin* **30** (1): 53–62. <https://doi.org/10.1023/A:1022067527584>
- Chernova, O. F. & Tselikova, T. N. 2004. *An atlas of mammalian hair: Fine structure of overhair and hair using scanning electron microscopy*. KMK, Scientific Press, Moscow, 430 pp. [In Russian]
- Chernova, O. F. 2006. Evolutionary aspects of hair polymorphism. *Biology Bulletin, Zoology* **33**(1): 43–52. <https://doi.org/10.1134/S1062359006010067>
- Chernova, O. F., Kirillova, I. V., Shapiro, B., Shidlovskiy, F. K., Soares, A. E. R., Levchenko, V. A. & Bertuch, F. 2016. Morphological and genetic identification and isotopic study of the hair of a cave lion (*Panthera spelaea* Goldfuss, 1810) from the Malyy Anyui River (Chukotka, Russia). *Quaternary Science Reviews* **142**: 61–73. <https://doi.org/10.1016/j.quascirev.2016.04.018>
- Chętnicki, W., Fedyk, S. & Bajkowska, U. 2007. Cases of coat colour anomalies in the Common Shrew, *Sorex araneus* L. *Folia biologica (Kraków)* **55**(1–2): 73–76. <https://doi.org/10.3409/173491607780006308>
- Clack, A. A., MacPhee, R. D. E. & Poinar, H. N. 2012. *Case study: Ancient sloth DNA recovered from hairs preserved in paleofeces*. In: Shapiro, B. & Hofreiter, M. (eds): *Ancient DNA: Methods and Protocols, Methods in Molecular Biology*, vol. 840; pp. 51–56. https://doi.org/10.1007/978-1-61779-516-9_1
- Cody, M. L. 1991. Honeyeater plucks koala for nest material. *Emu* **91**: 125–126. <http://dx.doi.org/10.1071/MU9910125>
- Cole, H. I. 1924. Taxonomic value of hair in Chiroptera. *Philippine Journal of Science, Manila* **24**: 117–121.
- Cooper, J. E & Cooper, M. E. 2013. *Wildlife forensic investigation: principles and practice*. CRC Press, 770 pp. <https://doi.org/10.1201/b14553>
- Csorba, G., Krivek, G., Sendula, T., Homonnay, Z. G., Hegyeli, Zs., Sugár, Sz., Farkas, J., Stojnic, N. & Németh, A. 2015. How can scientific researches change conservation priorities? A review of decade-long research of blind mole-rats (Rodentia: Spalacinae) in the Carpathian Basin. *Therya* **6**: 103–121. <https://doi.org/10.12933/therya-15-245>
- Czech, N. U., Klauer, G., Dehnhardt, G., & Siemers, B. M. 2008. Fringe for foraging? Histology of the bristle-like hairs on the tail membrane of the gleaning bat, *Myotis nattereri*. *Acta Chiropterologica* **10**(2): 303–311. <https://doi.org/10.3161/150811008X414872>
- Day, M. G. 1966. Identification of hair and feather remains in the gut and faeces of stoats and weasels. *Journal of Zoology, London* **148**: 201–217. <https://doi.org/10.1111/j.1469-7998.1966.tb02948.x>
- De, J. K. 1993. Study of surface structure of hair of some primates of Indian subcontinent. *Records of the Zoological Survey of India* **93**: 31–34.
- De, J. K., Chakraborty, S. & Chakraborty, R. 1998. Identification of dorsal guard hairs of five Indian species of Mongoose, *Herpestes* Illiger (Mammalia: Carnivora). *Mammalia* **62**: 285–295. <https://doi.org/10.1515/mamm.1998.62.2.285>

- Dearborn, N. 1939. Sections aid in identifying hair. *Journal of Mammalogy* **20**: 346–348. <https://doi.org/10.2307/1374262>
- Debelica, A. & Thies, M. L. 2009. *Atlas and key to the hair of terrestrial Texas mammals*. Special Publications, Museum of Texas Tech University, No 55; 102 pp. <https://doi.org/10.5962/bhl.title.142652>
- Debrot, S., Fivaz, G., Mermod, C. & Weber, J. M. 1982. *Atlas des Poils de Mammifères d'Europe*. Université de Neuchatel, 208 pp.
- De Marinis, A. M. & Agnelli, P. 1993. Guide to microscope analysis of Italian mammal hairs: Insectivora, Rodentia and Lagomorpha. *Bollettino di Zoologia* **60**: 225–232. <https://doi.org/10.1080/11250009309355815>
- De Marinis, A. M. & Asprea, A. 2006. Hair identification key of wild and domestic ungulates from southern Europe. *Wildlife Biology* **12**(3): 305–320. [https://doi.org/10.2981/0909-6396\(2006\)12\[305:HIKOWA\]2.0.CO;2](https://doi.org/10.2981/0909-6396(2006)12[305:HIKOWA]2.0.CO;2)
- Dhouailly, D. 2009. A new scenario for the evolutionary origin of hair, feather, and avian scales. *Journal of Anatomy* **214**: 587–606. <https://doi.org/10.1111/j.1469-7580.2008.01041.x>
- Dickman, C. R. 1986. A method for censusing small mammals in urban habitats. *Journal of Zoology, London* **210**: 631–636. <https://doi.org/10.1111/j.1469-7998.1986.tb03665.x>
- Dove, C. J. & Peurach, S. C. 2002. Microscopic analysis of feather and hair fragments associated with human mummified remains from Kagamil Island, Alaska. *Ethnographical Series, Department of Ethnography, The National Museum of Denmark, Copenhagen* **20**: 51–62.
- Ducommun, M., Jeanmaire-Besancon, F. & Vogel, P. (1994). Shield morphology of curly overhair in 22 genera of Soricidae (Insectivora, Mammalia). *Revue de Suisse Zoologie* **101**: 623–643. <https://doi.org/10.5962/bhl.part.79920>
- Dziurdzik, B. & Nowogrodzka-Zagórska, M. 1991. The histological structure of hairs of the giant panda, *Ailuropoda melanoleuca* (David, 1869), and the lesser panda, *Ailurus fulgens* (F. Cuvier, 1825), and the systematic position of these species. *Acta Zoologica Cracoviensia* **34**(2): 463–474.
- Dziurdzik, B. 1973. Key to the identification of hairs of mammals from Poland. *Acta Zoologica Cracoviensia* **18**: 73–92. [In Polish]
- Éhik, Gy. 1931. *Prémek és prémes állatok*. Szent István Társulat, Az Apostoli Szentszék Könyvkiadója, 152 pp. [in Hungarian]
- Fang, Y.-P., Lee, L.-L., Yew, F.-H. & Yu, H.-T. 1997. Systematics of white-toothed shrews (*Crocidura*) (Mammalia: Insectivora: Soricidae) of Taiwan: Karyological and morphological Studies. *Journal of Zoology* **242**: 151–166. <https://doi.org/10.1111/j.1469-7998.1997.tb02936.x>
- Feldhamer, G. A., Drickamer, L. C., Vessey, S. H., Merritt, J. F. & Krajewski, C. 2015. *Mammalogy: Adaptation, diversity, ecology*. Fourth edition. John Hopkins University Press, 394 pp.
- Fish, F. E. 2000. Biomechanics and energetics in aquatic and semiaquatic mammals: platypus to whale. *Physiological and Biochemical Zoology* **73**(6): 683–698. <https://doi.org/10.1086/318108>
- Fish, F. E., Smelstoys, J., Baudinette, R. V. & Reynolds, P. S. 2002. Fur does not fly, it floats: buoyancy of pelage in semi-aquatic mammals. *Aquatic Mammals* **28**(2): 103–112.
- Flesch, P. 1968. Epidermal iron pigments in red species. *Nature* **217**: 1056–1057. <https://doi.org/10.1038/2171056a0>

- Foran, D. R., Crooks, C. K. & Minta, S. C. 1997. DNA-based analysis of hair to identify species and individuals for population research and monitoring. *Wildlife Society Bulletin* **25**(4): 840–847.
- Friedenthal, H. 1911. *Tierhaaratlas*. Jena, Verlag von Gustav Fisher. 19 pp + 35 plates.
- Galatík, A., Galatík, J. Krul, Z. & Galatík, A. Jr. 2011. *FurSkin identification* <http://www.furskin.cz/about.htm>
- Galván, I., Garrido-Fernández, J., Ríos, J., Pérez-Gálvez, A., Rodríguez-Herrera, B. & Negro, J. J. 2016. Tropical bat as mammalian model for skin carotenoid metabolism. *Proceedings of the National Academy of Sciences of the United States of America (PNAS)* **113**: 10932–10937. <https://doi.org/10.1073/PNAS.1609724113>
- Garel, M., Cugnasse, J.-M., Gaillard, J. -M., Loison, A., Gibert, P., Douvre, P. & Dubray, D. 2005. Reproductive output of female mouflon (*Ovis gmelini musimon* × *Ovis* sp.): a comparative analysis. *Journal of Zoology, London* **266** (1): 65–71. <https://doi.org/10.1017/S0952836905006667>
- Gaubert, Ph., Wetz, M. & Chalubert, A. 2008. Genets and oyans interactive identification key. <http://lis.snv.jussieu.fr/genettes/web/>
- Gessner, C. 1551–1558. *Historiae animalium*. 4 vol. Zurich. Christoph Froschauer.
- Gessner, C. 1554. *Appendix historiae quadrupedum viviparorum & ouiparorum Conradi Gesneri Tigurini*. Excudebat C. Froschouerus. <http://dx.doi.org/10.5962/bhl.title.68305>
- Geyfman, M., & Andersen, B. 2010. Clock genes, hair growth and aging. *Aging (Albany NY)* **2**(3): 122–128. <https://doi.org/10.18632/aging.100130>
- Gharu, J. & Trevedi, S. 2016. Ancient hairs: need for morphological analysis of prehistoric and extant Mammals. *Vertebrate Zoology* **66**(2): 221–224.
- Gilmore, D. P., Da Costa, C. P. & Duarte, D. P. F. 2001. Sloth biology: an update on their physiological ecology, behavior and role as vectors of arthropods and arboviruses. *Brazilian Journal of Medical and Biological Research* **34**: 9–25. <https://doi.org/10.1590/S0100-879X2001000100002>
- Gloger, C. L. 1833. *Das Abändern der Vögel durch Einfluss des Klimas*. Breslau, Germany. <https://doi.org/10.5962/bhl.title.50395>
- González-Esteban, J., Villate, I. & Irizar, I. 2006. Differentiating hair samples of the European mink (*Mustela lutreola*), the American mink (*Mustela vison*) and the European polecat (*Mustela putorius*) using light microscopy. *Journal of Zoology* **270**: 458–461. <https://doi.org/10.1111/j.1469-7998.2006.00164.x>
- Grassé, P. P. 1955. *Mammifères. Fascicule II. Les ordres: Anatomie, Ethologie, Systématique*. Traité de zoologie, anatomie, systématique, biologie. Tome XVII. Masson et Cie. Paris, pp. 1173–2300.
- Gray, H. 1918. *Anatomy of the human body*. Lea & Febiger, Philadelphia & New York. pp. 1068–1069. <https://doi.org/10.5962/bhl.title.20311>
- Gregorin, R. & Cirranello, A. 2016. Phylogeny of Molossididae Gervais (Mammalia: Chiroptera) inferred by morphological data. *Cladistics* **32**: 2–35. <https://doi.org/10.1111/cla.12117>
- Griffiths, P. J. 2000. The evolution of feathers from dinosaur hair. *Gaia* **15**: 399–403.
- Grimm, J. & Waldeyer, W. 1884. *Atlas der menschlichen und tierischen Haare sowie der ähnlichen Fasergebilde: für die Bedürfnisse der Staatsarzneikunde, des Handels, der Technik und der Landwirtschaft*. Lahr, Druck und Verlag von Moritz Schauenburg. 195 pp.

- Grimmberger, E. 2017. *Die Säugetiere Mitteleuropas. Beobachten und Bestimmen*. Quelle & Meyer Verlag Wiebelsheim, 695 pp.
- Groves, C. P. 1994. *Morphology, habitat and taxonomy*. In: Boyd, L. & Houpt, K. (eds): Przewalski's horse. The history and biology of an endangered species. State University of New York Press, Albany, NY, p. 39–45.
- Gurnell, J., Lurz, P. W. W., Shirley, M. D. F., Cartmel, S., Garson, P. J., Magris, L. & Steele, J. 2004. Monitoring red squirrels *Sciurus vulgaris* and grey squirrels *Sciurus carolinensis* in Britain. *Mammal Review* **34**: 51–74. <https://doi.org/10.1046/j.0305-1838.2003.00028.x>
- Harničárová, K. & Adamík, P. 2016. Mammal hair in nests of four cavity-nesting songbirds: occurrence, diversity and seasonality. *Bird Study* **63**(2): 181–186. <http://dx.doi.org/10.1080/00063657.2016.1183584>
- Hausman, L. A. 1920. Structural characteristics of the hair of mammals. *The American Naturalist* **54** (635): 496–523. <https://doi.org/10.1086/279782>
- Hausman, L. A. 1920. A micrological investigation of the hair structure of the Monotremata. *American Journal of Anatomy* **27**(4): 463–495. <https://doi.org/10.1002/aja.1000270404>
- Hausman, L. A. 1920. The microscopic identification of commercial fur hairs. *Scientific Monthly* **10**: 7078.
- Hausman, L. A. 1920. Hairs that make fabrics. The microscopic identification of mammal hairs used in the textile industry. *The Scientific American* **1920**(2): 184, 200, 202.
- Hausman, L. A. 1924. Further studies on the relationships of the structural characters of mammalian hair. *American Naturalist* **58**: 544–557. <https://doi.org/10.1086/280006>
- Hausman, L. A. 1930. Recent studies of hair structure relationships. *Scientific Monthly* **30**(9): 258–277.
- Hausman, L. A. 1932. The cortical fuse of mammalian hair shafts. *American Naturalist* **66**: 461–470. <https://doi.org/10.1086/280452>
- Hausser, J. (ed.) 1995. *Säugetiere der Schweiz. Verbreitung, Biologie, Ökologie*. Birkhäuser Verlag, Basel, 501 pp.
- Hegyeli, Zs. & Kecskés, A. 2014. The occurrence of wild-living American Mink *Neovison vison* in Transylvania, Romania. *Small Carnivore Conservation* **51**: 23–28.
- Helgen, K. & Turvey, S. T. 2016. *Neovison macrodon*. The IUCN Red List of Threatened Species 2016: e. T40784A45204492. <https://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T40784A45204492.en>
- Henry, P. & Russello, M. A. 2011. Obtaining high quality DNA from elusive small mammals using low-tech hair snares. *European Journal of Wildlife Research* **57**(3): 429–435. <https://doi.org/10.1007/s10344-010-0449-y>
- Hess, W. M., Flinders, J. T., Pritchett, C. L. & Allen, J. V. 1985. Characterization of hair morphology in families Tayassuidae and Suidae with scanning electron microscopy. *Journal of Mammalogy* **66**(1): 75–84. <https://doi.org/10.2307/1380958>
- Heyn, A. N. J. 1954. *Fiber microscopy: a textbook and laboratory manual*. Interscience Publishers, Inc New York; 407 pp. <https://babel.hathitrust.org/cgi/pt?id=mdp.39015058432892;view=1up;seq=9>
- Hickey, M. B. C. & Fenton, M. B. 1987. Scent-dispersing hairs (osmetrichia) in some Pteropodidae and Molossidae (Chiroptera). *Journal of Mammalogy* **68**: 381–384. <https://doi.org/10.2307/1381478>

- Higginbotham, S., Wong, W. R., Linington, R. G., Spadafora, C., Iturrado, L. & Arnold, A. E. 2014. Sloth hair as a novel source of fungi with potent anti-parasitic, anti-cancer and anti-bacterial bioactivity. *PLoS ONE* **9**(1): e84549. <https://doi.org/10.1371/journal.pone.0084549>
- Higuchi, R., Bowman, B., Freiburger, M., Ryder, O. A. & Wilson, A. C. 1984. DNA-sequences from the Quagga, an extinct member of the horse family. *Nature* **312**(5991): 282–284. <https://doi.org/10.1038/312282a0>
- Hill, P., Brantley, H. & Van Dyke, M. 2010. Some properties of keratin biomaterials: Kerateines. *Biomaterials* **31**: 585–593. <https://doi.org/10.1016/j.biomaterials.2009.09.076>
- Hoey, K., Wise, R., & Adler, G. 2004. Ultrastructure of echimyid and murid rodent spines. *Journal of Zoology* **263**(3): 307–315. <https://doi.org/10.1017/S0952836904005370>
- Hoffmann, H. J. 2000. Messel: Window on an ancient world. *National Geographic Magazine* **197**(2): 34–51.
- Hollemeier, K., Altmeyer, W., & Heinzle, E. 2002. Identification and quantification of feathers, down and hair of avian and mammalian origin using MALDI-TOF-mass spectrometry. *Analytical Chemistry* **74**: 5960–5968. <https://doi.org/10.1021/ac020347f>
- Hollemeier K., Altmeyer W. & Heinzle E. 2007. Identification of furs of domestic dog, raccoon dog, rabbit and domestic cat by hair analysis using MALDI-TOF mass spectrometry. *Spectroscopy Europe* **19**(2): 8–15.
- Homan, J. A. & Genoways, H. H. 1978. An analysis of hair structure and its phylogenetic implications among Heteromyd Rodents. *Journal of Mammalogy* **59**(4): 740–760. <https://doi.org/10.2307/1380139>
- Hooke, R. 1665. *Micrographia, or some physiological descriptions of minute bodies made by magnifying glasses with observations and inquiries thereupon*. J. Martyn and J. Allestry, Printers to the Royal Society, London, 323 pp.
- Houck, M. M. & Budowle, B. 2002. Correlation of microscopic and mitochondrial DNA hair comparisons, *Journal of Forensic Sciences* **47**: 964–967. <https://doi.org/10.1520/JFS15515J>
- Hutterer, R. & Hürter, T. 1981. Adaptive Haarstrukturen bei Wasserspitzmäusen (Insectivora, Soricinae). *Zeitschrift für Saugetierkunde* **46**: 1–11.
- Ibara, B., Itzel, I. & Sánchez-Cordero, V. 2004. Catálogo de pelos de guardia dorsal en mamíferos del estado de Oaxaca, México. *Anales del Instituto de Biología. Serie Zoología, Universidad Nacional Autónoma de México* **75**(2): 383–437.
- Imes, D. L., Geary, L. A., Grahn, R. A. & Lyons, L. A. 2006. Albinism in the domestic cat (*Felis catus*) is associated with a tyrosinase (TYR) mutations. *Animal Genetics* **37**: 175–178. <https://doi.org/10.1111/j.1365-2052.2005.01409.x>
- Ji, Q., Luo, Z. -X., Yuan, C. -X. & Tabrum, A. R. 2006. A swimming mammaliaform from the Middle Jurassic and ecomorphological diversification of early mammals. *Science* **311**(5764): 1123–1127. <https://doi.org/10.1126/science.1123026>
- Ji, Q., Luo, Z. -X., Yuan, C. -X., Wible, J. R., Zhang, J. -P. & Georgi, J. A. 2002. The earliest known eutherian mammal. *Nature* **416**(6883): 816–822. <https://doi.org/10.1038/416816a>
- Keller, A. 1981. Détermination des mammifères de la Suisse par leur pelage: V. Carnivora, VI. Artiodactyla. *Revue Suisse de Zoologie* **88**(3): 803–820. <https://doi.org/10.5962/bhl.part.82410>

- Keller, A. 1984. Etude de la structure fines des jarres dorsaux de quelques Canidés sauvages et domestiques du genre *Canis* (Mammalia: Canidae). *Revue Suisse de Zoologie* **91**: 973–992. <https://doi.org/10.5962/bhl.part.81593>
- Keller, A. 1992. Notes sur une étude comparative des jarres primaires de trois espèces d'Equidae: *Equus asinus*, *E. przewalskii* et *E. caballus*. *Revue Suisse de Zoologie* **99**(4): 735–739. <https://doi.org/10.5962/bhl.part.79849>
- Kelly, M. J. 2001. Computer-aided photograph matching in studies using individual identification: an example from Serengeti cheetahs. *Journal of Mammalogy* **82**(2): 440–449. <https://doi.org/10.5962/bhl.part.79849>
- Kenagy, G. J. & Pearson, O. P. 2000. Life with fur and without: experimental field energetics and survival of naked meadow voles. *Oecologia* **122**(2): 220–224. <https://doi.org/10.1007/PL00008849>
- Kendall, K. C. & McKelvey, K. S. 2008. *Hair collection*. In: Long, R. A., MacKay, P. Zielinski, W. J. & Ray, J. C. (eds): *Noninvasive survey methods for Carnivores 2008*. Island Press, Washington, D. C., p. 135–176.
- Kennedy, A. J. 1982. Distinguishing characteristics of the hairs of wild and domestic canids from Alberta. *Canadian Journal of Zoology* **60**(4): 536–541. <https://doi.org/10.1139/z82-080>
- Keogh, H. J. 1985. A photographic reference system based on the cuticular scale patterns and groove of the hair of 44 species of Southern African Cricetidae and Muridae. *South African Journal of Wildlife Research*, Stellenbosch **15**: 109–159.
- Kingdon, J., Agwanda, B., Kinnaird, M., O'Brien, T., Holland, C., Gheysens, T., Boulet-Audet, M. & Vollrath, F. 2011. A poisonous surprise under the coat of the African crested rat. *Proceedings of the Royal Society B, Biological Sciences* **279**(729): 675–680. <https://doi.org/10.1098/rspb.2011.1169>
- Kirk, P. L. 1940. Human hair studies: 1. General considerations of hair individualization and its forensic importance. *Journal of Criminal Law and Criminology* **31**: 486–496. <https://doi.org/10.2307/1137587>
- Koon, D. W. 1998. "Is Polar Bear Hair Fiber Optic?". *Applied Optics* **37**: 3198–3200. <https://doi.org/10.1364/AO.37.003198>
- Koppiker, B. R. & Sabnis J. H. 1976. Identification of hairs of some Indian mammals. *Journal of the Bombay Natural History Society* **73**: 5–20.
- Krstić, R. V. 1997. *Human microscopic anatomy*. Springer Verlag, Berlin, Heidelberg, New York, Tokyo, pp. 450–459.
- Kryštufek, B. & Vohralík, V. 2012. Taxonomic revision of the Palearctic rodents (Rodentia). Sciuridae: Xerinae 1 (*Eutamias* and *Spermophilus*). *Lynx* **43**(1–2): 17–111.
- Kuhn, R. A., Ansoerge, H., Godynicki, S., & Meyer, W. 2010. Hair density in the Eurasian otter *Lutra lutra* and the Sea otter *Enhydra lutris*. *Acta Theriologica* **55**(3): 211–222. <https://doi.org/10.4098/j.at.0001-7051.014.2009>
- Kuhn, R. A. & Meyer, W. 2010. Comparative hair structure in the Lutrinae (Carnivora: Mustelidae). *Mammalia*. **74**: 291–303. <https://doi.org/10.1515/MAMM.2010.039>
- Kurihara, N. 2013. Mitochondrial DNA in hair shafts from study skin. *Genetics and Molecular Research* **12**(4): 5396–5404. <https://doi.org/10.4238/2013.November.11.1>

- Lambert, M. & Balthazard, V. 1910. *Le poil de l'homme et des animaux. Applications aux expertises médico-légales et aux expertises des fourrures*. Paris, Steinheil G. éditeur, 2, Rue Casimir-Delavigne. 228 pp. (plus Planches XXXIV.)
- Lapini, L. & Testone, R. 1998. A new *Sorex* from north-eastern Italy (Mammalia: Insectivora: Soricidae). *Gortania Atti Museo Friul di Storia Naturale* **20**: 233–252.
- Lee, E., Choi, T.-Y., Woo, D., Min, Mi.-S., Sugita, S. & Lee, H. 2014. Species identification key of Korean Mammal Hair. *Journal of Veterinary Medical Science, Wildlife Science* **76**(5): 667–675. <https://doi.org/10.1292/jvms.13-0569>
- Leeuwenhoek Mr. 1665–1678. Microscopical observations from Mr. Leeuwenhoeck, about blood, milk, bones, the brain, spittle, cuticula, sweat, fatt, teares; communicated in two letters to the publisher. *Philosophical Transactions* **1674** (1665–1678) **9**: 121–131.
- Liwanag, H. E. M., Berta, A., Costa, D. P., Abney, M. & Williams, R. M. 2012. Morphological and thermal properties of mammalian insulation: the evolution of fur for aquatic living. *Biological Journal of the Linnean Society* **106**: 926–939. <https://doi.org/10.1111/j.1095-8312.2012.01900.x>
- Lochte, Th. 1938. *Atlas der Menschlichen und Tierischen Haare*. Verlag Dr. Paul Schöps, Leipzig, 306 pp.
- Lochte, Th. 1954. *Tafeln zur Haarkunde, Beiträge zur Haut-, Haar- und Fellkunde*. Leipzig, vol. 6.
- López-Baucells, A., Mas, M., Puig-Montserrat, X. & Flaquer, C. 2013. Hipo-pigmentation in vespertilionid bats: first record of a leucistic soprano pipistrelle (*Pipistrellus pygmaeus*). *Barbastella* **6**(1): 63–70.
- Lucati, F. & López-Baucells, A. 2017. Chromatic disorders in bats: a review of pigmentation anomalies and the misuse of terms to describe them. *Mammal Review* **47**(2): 112–123. <https://doi.org/10.1111/mam.12083>
- Lyne, A. G. & McMahon, T. S. 1951. Observations on the surfaces the hairs of Tasmanian monotremes and marsupials. *Papers and Proceedings of the Royal Society of Tasmania* **1950**: 71–84.
- MacDaniel, G. W. 2000. Efficacy of lures and hair snares to detect *Lynx*. *Wildlife Society Bulletin* **28**: 119–123.
- Maderson, P. F. A. 1972. When? Why? And How? Some speculations on the evolution of the Vertebrate integument. *American Zoologist* **12**: 159–171. <https://doi.org/10.1093/icb/12.1.159>
- Maderson, P. F. A. 2003. Mammalian skin evolution: a re-evaluation. *Experidermal Dermatology* **12**: 233–236. <https://doi.org/10.1034/j.1600-0625.2003.00069.x>
- Martin, T., Marugán-Lobón, J., Vullo, R., Martín-Abad, H., Luo, Z. -X. & Buscalioni, A. D. 2015. A Cretaceous eutriconodont and integument evolution in early mammals. *Nature* **526**(7573): 380. <https://doi.org/10.1038/nature14905>
- Mathiak, H. A. 1938a. A rapid method of cross-sectioning mammalian hairs. *Journal of Wildlife Management* **2**: 162–164. <https://doi.org/10.2307/3796437>
- Mathiak, H. A. 1938b. Key to mammalian hairs found in faeces. *Journal of Wildlife Management* **2**: 251–268. <https://doi.org/10.2307/3795673>

- Mayer, W. V. 1952. The hair of California mammals with keys to the dorsal guard hairs of California mammals. *American Midland Naturalist* **48**: 480–512. <https://doi.org/10.2307/2422262>
- Mckittrick, J., Chen, P. Y., Bodde, S. G., Yang, W., Novitskaya, E. E., & Meyers, M. A. 2012. The structure, functions, and mechanical properties of keratin. *Journal of the Minerals, Metals & Materials Society* **64**(4): 449–468. <https://doi.org/10.1007/s11837-012-0302-8>
- Meng, J. & Wyss, A. R. 1997. Multituberculate and other mammal hair recovered from Paleogene excreta. *Nature* **385**: 712–714. <https://doi.org/10.1038/385712a0>
- Meng, J., Hu, Y., Wang, Y., Wang, X. & Li, C. 2006. A Mesozoic gliding mammal from northeastern China. *Nature* **444**(7121): 889–893. <https://doi.org/10.1038/nature05234>
- Meyer, M., Palkopoulou, E., Baleka, S., Stiller, M., Penkman, K. E. H., Alt, K. W., Ishida, Y., Mania, D., Mallick, S., Meijer, T., Meller, H., Nagel, S., Nickel, B., Ostritz, S., Rohland, N., Schauer, K., Schüler, T., Roca, A. L., Reich, D., Shapiro, B., Hofreiter, M. 2017. Palaeogenomes of Eurasian straight-tusked elephants challenge the current view of elephant evolution. *eLife* **6**: e25413. <https://doi.org/10.7554/eLife.25413>
- Meyer, W., Hülmann, G. & Seger, H. 2002. *SEM-Atlas of the hair cuticle structure of central European mammals*. Verlag M. & H. Shaper Alfeld, Hannover, 248 pp.
- Meyer, W., Hülmann, G. & Seger, H. 1995. Remarks on specific adaptive scale structure of the hair cuticle in some European bats. *European Journal of Morphology* **33**: 509–513.
- Meyer, W. A., Schnapper, A., Hülmann, G. & Seger, H. 2000. Domestication-related variations of the hair cuticula pattern in mammals. *Journal of Animal Breeding and Genetics* **117**: 281–283. <https://doi.org/10.1111/j.1439-0388.2000.00260.x>
- Miles, W. B. 1965. Studies of the cuticular structure of the hairs of Kansas bats. *Search* **5**: 48–50.
- Miller-Butterworth, C. M., Murphy, W. J., O'Brien, S. J., Jacobs, D. S., Springer, M. S. & Teeling, E. C. 2004. A family matter: conclusive resolution of the taxonomic position of the Long-Fingered Bats, *Miniopterus*. *Molecular Biology and Evolution* **24**(7): 1553–1561. <https://doi.org/10.1093/molbev/msm076>
- Mitchell-Jones, A. J., Amori, G., Bogdanowicz, W., Kryštufek, B., Reijnders, P. J. H., Spitzenberger, F., Stubbe, M., Thissen, J. B. M., Vohralk, V. & Zima, J., 1999. *The atlas of European mammals*. Academic Press, London, 496 pp.
- Moore, T. D., Spence, L. E. & Dugnolle, C. E. 1974. Identification of the dorsal guard hairs of some mammals of Wyoming. *Wyoming Game and Fish Department Bulletin* **14**, 177 pp.
- Morris, P. A. & Tutt, A. 1996. Leucistic hedgehogs on the island of Alderney. *Journal of Zoology* **239**: 387–389. <https://doi.org/10.1111/j.1469-7998.1996.tb05458.x>
- Mowat, G. & Paetkau, D. 2002. Estimating marten *Martes americana* population size using hair capture and genetic tagging. *Wildlife Biology* **8**(3): 201–209.
- Mowat, G. & Strobeck, C. 2000. Estimating population size of grizzly bears using hair capture, DNA profiling and mark-recapture analysis. *Journal of Wildlife Management* **64**: 183–193. <https://doi.org/10.2307/3802989>
- Müller-Schwarze, D., Volkman, N. J. & Zemanek, K. F. 1977. Osmetricchia: specialized scent hair in black-tailed deer. *Journal of Ultrastructure Research* **59**: 223–230. [https://doi.org/10.1016/S0022-5320\(77\)90035-1](https://doi.org/10.1016/S0022-5320(77)90035-1)

- Nason, E. S. 1948. Morphology on hair of eastern North American bats. *American Midland Naturalist* **39**(2): 345–361. <https://doi.org/10.2307/2421589>
- Nedyalkov, N., Koshev, Y., Raykov, I. & Bardarov, G. 2014. Color variation of small mammals' (Mammalia: Rodentia and Insectivora) coats from Bulgaria. *North-Western Journal of Zoology* **10**(2): 314–317.
- Newman, C., Buesching, C. D. & Wolf, J. O. 2005. The function of facial masks in “midguild” carnivores. *Oikos* **108**: 623–633. <https://doi.org/10.1111/j.0030-1299.2005.13399.x>
- Nielsen, K. P., Zhao, L., Stamnes, J. J., Stamnes, K., & Moan, J. 2008. The optics of human skin: Aspects important for human health. *Solar Radiation and Human Health. The Norwegian Academy of Science and Letters*, pp 35–46.
- Noback, C. R. 1951. Morphology and phylogeny of hair. *Annales of the New York Academy Science* **53**: 476–492. <https://doi.org/10.1111/j.1749-6632.1951.tb31950.x>
- Nowak, R. M. & Paradiso, J. L. 1983. *Walker's Mammals of the World*. 4th edition. Johns Hopkins University Press, Baltimore, 1362 pp.
- Ogden, R., Dawney, N. & McEwing, R. 2009. Wildlife DNA forensic – bridging the gap between conservation genetics and law enforcement. *Endangered Species Research* **9**: 179–195. <https://doi.org/10.3354/esr00144>
- Ondrusova, K., & Adamík, P. 2013. Characterizing the mammalian hair present in Great Tit (*Parus major*) nests. *Bird Study* **60** (3): 428–431. <https://doi.org/10.1080/00063657.2013.818935>
- Ortolani, A. 1998. Spots, stripes, tail trips and dark eyes: Predicting the function of carnivore colour patterns using the comparative method. *Biological Journal of the Linnean Society* **67**: 433–476. <https://doi.org/10.1111/j.1095-8312.1999.tb01942.x>
- O'Sullivan, N. J., Teasdale, M. D., Mattiangeli, V., Maixner, F., Pinhasi, R., Bradley, D. G. & Zink, A. 2016. A whole mitochondria analysis of the Tyrolean Iceman's leather provides insights into the animal sources of Copper Age clothing. *Scientific Reports* **6**, 31279. <https://doi.org/10.1038/srep31279>
- Oyer, E. R. 1946. Identification of mammals from studies of hair structure. *Transactions of the Kansas Academy of Science* (1903) **49**(2): 155–160. <https://doi.org/10.2307/3625853>
- Parakkal, P. F. & Alexander, N. J. 1972. *Keratinization. A survey of vertebrate epithelia*. Academy Press, New York, London, 59 pp.
- Patkó, L., Ujhegyi, N., Szabó, L., Péter, F., Schally, G., Tóth, M., Lanszki, J., Nagy, Z., Szemethy, L., & Heltai, M. 2015. Even a hair casts its shadow: review and testing of noninvasive hair collecting methods of carnivore species. *North-Western Journal of Zoology* **12**(1): 122–129.
- Patterson, B. D. & Norris, R. W. 2016. Towards a uniform nomenclature for ground squirrels: the status of the Holarctic chipmunks. *Mammalia* **80**(3): 241–25. <https://doi.org/10.1515/mammalia-2015-0004>
- Pech-Caché, J. M., Sosa-Escalante, J. E. & Koyoz Cruz, M. E. 2009. Guía para la identificación de pelos de guardia de Mamíferos no voladores del estado de Yucatán, México. *Revista Mexicana de Mastozoología* **13**: 7–33.
- Peñalver, E. & Grimaldi, D. 2006. Assemblages of mammalian hair and blood-feeding midges (Insecta: Diptera: Psychodidae: Phlebotominae) in Miocene Amber. *Transactions of*

- the Royal Society of Edinburgh, Earth Sciences* **96**: 177–195. <https://doi.org/10.1017/S0263593300001292>
- Petter, J. J. & Petter-Rousseaux, A. 1963. Notes biologiques sur les Centetinae. *La Terre et la Vie* **17**: 66–80.
- Pierallini, R., Keller, A. & Moretti, M. 2004. Chiave di determinazione dei Chiroterteri (Mammalia) della Svizzera attraverso l'osservazione al microscopio ottico della struttura dei peli. *Revue Suisse de Zoologie* **111**(2): 381–393. <https://doi.org/10.5962/bhl.part.80244>
- Po-Chedly, D. S. 1951. A method for staining the scales of coarse hairs. *Journal of Mammalogy* **32**: 232. <https://doi.org/10.1093/jmammal/32.2.232>
- Pocock, R. I. 1914. On the facial vibrissae of Mammalia. *Proceedings of the Zoological Society of London* **84**: 889–912. <https://doi.org/10.1111/j.1469-7998.1914.tb07067.x>
- Poinar, G. O. 1988. Hair in Dominican amber: Evidence for tertiary land mammals in the Antilles. *Experientia* **44**: 88–89. <https://doi.org/10.1007/BF01960261>
- Poulton, E. B. 1890. *The colours of animals, their meaning and use, especially considered in the case of insects*. London, Kegan Paul, Trench & Trübner. <https://doi.org/10.5962/bhl.title.69899>
- Quadros, J. & Monteiro-Filho, E. L. A. 1998. Effects of digestion, putrefaction, and taxidermy processes on *Didelphis albiventris* hair morphology. *Journal of Zoology, London* **244**: 331–334. <https://doi.org/10.1111/j.1469-7998.1998.tb00037.x>
- Quadros, J. & Monteiro-Filho, E. L. A. 2006. Revisão conceitual, padrões microestruturais e proposta nomenclatória para os pêlos-guarda de mamíferos brasileiros. *Revista Brasileira de Zoologia* **23**(1): 279–292. <https://doi.org/10.1590/S0101-81752006000100023>
- Quekett, J. 1844. On the structure of bat's hair. *Transactions of the Microscopical Society of London* **1**: 58–62. <https://doi.org/10.1111/j.1365-2818.1844.tb04618.x>
- Quick, D. E. & Hillenius, W. J. 2013. *Dinosaur physiology: were dinosaurs warm-blooded?* John Wiley & Sons Ltd, Chichester. <https://doi.org/10.1002/9780470015902.a0003323.pub2>
- Ray, J. 1693. Synopsis methodica animalium quadrupedum et serpentini generis: vulgarium notas características, rariorum descriptiones integras exhibens: cum historiis & observationibus anatomicis perquam curiosis: praemittuntur nonnulla De animalium in genere, sensu, generatione, divisione, &c. / Synopsis animalium quadrupedum. Londini. Impensis S. Smith & B. Walford. <http://dx.doi.org/10.5962/bhl.title.41459>
- Rensch, B. 1938. Some problems of geographical variation and species formation. *Proceedings of the Linnean Society of London* **150**: 275–285. <https://doi.org/10.1111/j.1095-8312.1938.tb00182k.x>
- Rose, K. D. & Gaudin, T. J. 2010. Xenarthra and Pholidota (Armadillos, Anteaters, Sloths and Pangolins). In: eLS. John Wiley & Sons Ltd, Chichester. <https://doi.org/10.1002/9780470015902.a0001556.pub2>
- Sahajpal, V., Goyal S. P., Thakar, M. K. & Jayapal, R. 2009. Microscopic hair characteristics of a few bovid species listed under Schedule-I of Wildlife (Protection) Act 1972 of India. *Forensic Science International* **189**: 34–45. <https://doi.org/10.1016/j.forsci-int.2009.04.008>
- Santana, S. E., Dial, T. O., Eiting, T. & Alfaro, M. E. 2011. Roosting ecology and the evolution of pelage markings in bats. *PLoS ONE* **6**(10): e25845. <https://doi.org/10.1371/journal.pone.025845>

- Sarko, D. K., Rice, F. L. & Reep, R. L. 2011. Mammalian tactile hair: divergence from a limited distribution. *Annals of the New York Academy of Sciences* **1225**: 90–100. <https://doi.org/10.1111/j.1749-6632.2011.05979.x>.
- Sato, J. J., Wolsan, M., Minami, S., Hosoda, T., Sinaga, M. H., Hiyama, K., Yamaguchi, Y. & Suzuki, H. 2009. Deciphering and dating the red panda's ancestry and early adaptive radiation of Musteloidea. *Molecular Phylogenetics and Evolution* **53**(3): 907–922. <https://doi.org/10.1016/j.ympev.2009.08.019>
- Scotts, D. J. & Craig, S. A. 1988. Improved hair-sampling tube for the detection of rare mammals. *Australian Wildlife Research* **15**: 469–472. <https://doi.org/10.1071/WR9880469>
- Sebetic, K., Sjerobabski-Masnec, I., Cavka, V., Biljan, D., Krolo, I. 2008. UV damage of the hair. *Collegium Antropologicum* **32**(2): 163–165.
- Seiler, N. 2010. SEM-Atlas of hair structures of South-African mammals. *Mammalia* **74**: 281–290. <https://doi.org/10.1515/mamm.2010.033>
- Sessions, B. D., Nielson, C. E., Sowa, J. M., Hess, W. M., Skidmore, W. & Carmack, B. A. 2014. Is bat hair morphology exceptional? *Vespertilio* **17**: 171–183.
- Shapiro, B. & Hofreiter, M. (eds) 2012. *Ancient DNA: methods and protocols, methods in molecular biology*, Vol. 840, 248 pp., Springer Science + Business Media, LLC 201 Humana Press, New York. https://doi.org/10.1007/978-1-61779-516-9_1
- Short, H. L. 1978. Analysis of cuticular scales on hair using the scanning electron microscope. *Journal of Mammalogy* **59**: 261–268. <https://doi.org/10.2307/1379911>
- Slominski, A., Tobin, D. J., Shibahara, S., & Wortsman, J. 2004. Melanin pigmentation in mammalian skin and its hormonal regulation. *Physiological Reviews* **84**(4): 1155–1228. <https://doi.org/10.1152/physrev.00044.2003>
- Snyder, H. K., Maia, R., D'Alba, L., Shultz, A. J., Rowe, K. M. C., Rowe, K. C., Shawkey, M. D. 2012. Iridescent colour production in hairs of blind golden moles (Chrysochloridae). *Biology Letters* **8**(3): 393–396. <https://doi.org/10.1098/rsbl.2011.1168>
- Sokolov, V. E. 1982: *Mammal skin*. University of California Press Ltd, 688 pp.
- Spearman, R. I. C. 1966. The keratinization of epidermal scales, feathers and hairs. *Biological Reviews* **41**: 59–96. <https://doi.org/10.1111/j.1469-185X.1966.tb01538.x>
- Spearman, R. I. C. 1967. On the nature of the horny scales of the pangolin. *Journal of the Linnean Society of London, Zoology* **46**: 267–273.
- Spencer, B. & Sweet, G. 1899. The Structure and Development of the Hairs of Monotremes and Marsupials. Part. 1. Monotremes. *The Quarterly Journal of Microscopical Science* **41**: 549–588.
- Spitzenberger, F. 2001. *Die Säugetierfauna Österreichs*. Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft, 895 pp.
- Stains, H. J. 1958. Field key to guard hair of middle western furbearers. *Journal of Wildlife Management* **22**(1): 95–97. <https://doi.org/10.2307/3797306>
- Stankowich, T., Caro, T. & Cox, M. 2011. Bold coloration and the evolution of aposematism in terrestrial carnivores. *Evolution* **65**(11): 3090–3099. <https://doi.org/10.1111/j.1558-5646.2011.01334>
- Stein, H. W. 1960. Zum Haarwechsel der Feldmaus, *Microtus arvalis* (Pallas, 1779) und weiterer Muroidea. *Acta Theriologica* **3**: 27–44. <https://doi.org/10.4098/AT.arch.60-3>

- Sterbing-D'Angelo, S. J., Chadha, M., Marshall, K. L. & Moss, C. F. 2017. Functional role of airflow-sensing hairs on the bat wing. *Journal of Neurophysiology* **117**: 705–712. <https://doi.org/10.1152/jn.00261.2016>
- Steyer, K., Simon, O., Kraus, R. H. S., Haase, P. & Nowak, C. 2012. Hair trapping with valeriana treated lure sticks as a tool for genetic wildcat monitoring in low-density habitats. *European Journal of Wildlife Research* **59**: 39–46. <https://doi.org/10.1007/s10344-012-0644-0>
- Stoner, C. J., Bininda-Emonds, O. R. P. & Caro, T. 2003. The adaptive significance of coloration in lagomorphs. *Biological Journal of the Linnean Society* **79**: 309–328. <https://doi.org/10.1046/j.1095-8312.2003.00190.x>
- Storch, G. 1992. The mammals of Island Europe. *Scientific American* **1992**(Feb.): 64–69. <https://doi.org/10.1038/scientificamerican0292-64>
- Suckling, G. C. 1978. A hair sampling tube for the detection of small mammals in trees. *Australian Wildlife Research* **5**: 249–252. <https://doi.org/10.1071/WR9780249>
- Swift J. A. & Smith J. R. 2001. Microscopical investigations on the epicuticle of mammalian keratin fibres. *Journal of Microscopy* **204**: 203–211. <https://doi.org/10.1046/j.1365-2818.2001.00957.x>
- Szunyogh, J. & Topál, Gy. 1972. *Emlősök – Mammalia. Rovarevők – Insectivora*. Magyarország Állatvilága [Fauna of Hungary] XXII. Budapest, 55 pp. [In Hungarian]
- Taru, P. & Backwell, L. R. 2013. Identification of fossil hairs in *Parahyaena brunnea* coprolites from Middle Pleistocene deposits at Gladysvale cave, South Africa. *Journal of Archaeological Science* **40**: 3674–3685. <https://doi.org/10.1016/j.jas.2013.04.031>
- Taru, P. & Backwell, L. R., 2014. Hair morphology of some artiodactyls from southern Africa. *Annals of the Ditsong National Museum of Natural History* **4**: 26–32.
- Taylor, R. J. 1985. Identification of the hair of Tasmanian mammals. *Papers and Proceedings of the Royal Society of Tasmania*, Hobart **119**: 69–82.
- Teerink, B. J. 1991. *Hair of West European Mammals: Atlas and identification key*. Cambridge University Press, Cambridge, 233 pp.
- Tobin, D. J. & Paus, R. 2001. Graying: gerontobiology of the hair follicle pigmentary unit. *Experimental Gerontology* **36**: 29–54. [https://doi.org/10.1016/S0531-5565\(00\)00210-2](https://doi.org/10.1016/S0531-5565(00)00210-2)
- Toldt, K, von. 1910. Über eine beachtenswerte Haarsorte und über das Haarform System der Säugetiere. *Annalen des K. K. Naturhistorischen Hofmuseums* **24**: 195–268.
- Toldt, K, von. 1935. *Aufbau und Natürliche Färbung des Haarkleides der Wildsäugetiere*. Deutsche Gesellschaft für Kleintier und Pelztierzucht GmbH & Co. Leipzig. 291 pp.
- Tóth, A. M. 2002. A szőrscapdázás alkalmazásának lehetőségei és korlátai. *Vadbiológia* **9**: 117–122. [in Hungarian]
- Tóth, A. M. 2002. Identification of Hungarian Mustelidae and other small carnivores using guard hair analysis. *Acta Zoologica Academiae Scientiarum Hungaricae* **48**(3): 237–250.
- Tóth, M. 2003. *Az emlősök szőrmintáinak információtartalma, a szőrhatózás módszertana és a módszer gyakorlati alkalmazása*. Doktori értekezés. Eötvös Loránd Tudományegyetem, Állattrendszertani és Ökológiai Tanszék. 141 pp. [in Hungarian]
- Tóth, M. 2008. A new non-invasive method for detecting mammals from birds' nests. *Journal of Wildlife Management* **72**(5): 1237–1240. <https://doi.org/10.2193/2006-501>

- Tóth, M. 2015. *A magyar emlősfauna szőrtani kézikönyve*. Magyar Természettudományi Múzeum, Budapest, 208 pp. [in Hungarian, with English subtitle: Handbook of trichomorphology of Hungarian mammals.]
- Tóth, M. & Heltai, M. 2010. Esettanulmányok. In: Heltai, M. (ed.): *Emlős ragadozók Magyarországon. Elterjedés, ökológiai szerep, gazdálkodás és védelem*. Mezőgazda Kiadó, Budapest, pp. 121–210. [in Hungarian]
- Tridico, S. R., Koch, S., Michaud, A., Thomson, G., Kirkbride, K. P. & Bunce, M. 2014. Interpreting biological degradative processes acting on mammalian hair in the living and the dead: which ones are taphonomic? *Proceedings of Biological Sciences* **281**(1796): 20141755. <http://dx.doi.org/10.1098/rspb.2014.1755>
- Tupinier, Y. 1973. Morphologie des poils de Chiroptères d'Europe occidentale par étude du microscope électronique à balayage. *Revue Suisse de Zoologie* **80**: 635–653. <https://doi.org/10.5962/bhl.part.75960>
- Ujhelyi, P. 1994. *A magyarországi vadonélő emlősállatok határozója*. MME, Budapest, 189 pp. [in Hungarian]
- Valente, A. 1983. Hair structure of the Woolly mammoth, *Mammuthus primigenius* and the modern elephants, *Elephas maximus* and *Loxodonta africana*. *Journal of Zoology, London* **199**: 271–274. <https://doi.org/10.1111/j.1469-7998.1983.tb02095.x>
- Valente, A. M., Gomes Rocha, R., Lozano, E., Ferreira, J. P., Fonseca, C. 2015. Atlas dos Pelos dos Mamíferos Terrestres Ibéricos. Edições afrontamento, Edição de coleção Fernando Morgado e Amadeu Soares, Coleção Biologicando (Atlas of hairs of Iberian terrestrial mammals.), 192 pp. <https://doi.org/10.1111/j.1469-7998.1983.tb02095.x>
- Velasco, M. V. R., Dias, T. C. S., Freitas, A. Z., Vieira Júnior, N. D., Pinto, C. A. S. O., Kaneko, T. M. & Baby, A. R. 2009. Hair fiber characteristics and methods to evaluate hair physical and mechanical properties. *Brazilian Journal of Pharmaceutical Sciences* (45)1: 153–162. <https://doi.org/10.1590/S1984-82502009000100019>
- Venkatraman, K., De, J. K. & Tandon, S. K. (1994): Ultrastructural studies of hair of seventeen Carnivores using scanning electron micrographs. *Records of the Zoological Survey of India* **94**: 145–158.
- Veron, G., Patou, M. -L., Tóth, M., Goonatilake, M. & Jennings, M. A. 2015. How many species of *Paradoxurus* civets are there? New insights from India and Sri Lanka. *Journal of Zoological Systematics and Evolutionary Research* **53**(2): 161–174. <https://doi.org/10.1111/jzs.12085>
- Vincent, J. F. V. & Owers, P. 1986. Mechanical design of hedgehog spines and porcupine squills. *Journal of Zoology, London* **210**: 55–75. <https://doi.org/10.1111/j.1469-7998.1986.tb03620.x>
- Vullo, R., Girard, V., Azar, D. & Néraudeau, D. 2010. Mammalian hairs in early Cretaceous amber. *Naturwissenschaften* **97**(7): 683–687. <https://doi.org/10.1007/s00114-010-0677-8>
- Wallace, A. R. 1877. The colors of animals and plants. *The American Naturalist* **11**(11): 641–662. <https://doi.org/10.1086/271979>
- Wallis, R. L. 1993. A key for the identification of guard hairs of some Ontario mammals. *Canadian Journal of Zoology* **71**: 587–591. <https://doi.org/10.1139/z93-080>

- Walter, W. D., Kurle, C. M. & Hopkins, J. B. 2014. Applications of stable isotope analysis in mammalian ecology. *Isotopes in Environmental and Health Studies* **50**(3): 287–290. <https://doi.org/10.1080/10256016.2014.933214>
- Wildman, A. & Manby, J. 1938. The hairs of the Monotremata, with special reference to their cuticular-scale pattern. *Transactions of the Royal Society of Edinburgh* **59**(2): 333–349. <https://doi.org/10.1017/S008045680000911X>
- Williams, C. S. 1938. Aids to the identification of mole and shrew hairs with general comments on hair structure and hair determination. *Journal of Wildlife Management* **2**: 239–250. <https://doi.org/10.2307/3795672>
- Williamson, V. H. H. 1951. Determination of hairs by impressions. *Journal of Mammalogy* **32**: 80–105. <https://doi.org/10.2307/1375415>
- Wilson, D. E. & Reeder, D. M. (eds) 2005. *Mammal species of the World. A taxonomic and geographic reference*. 3rd ed. Johns Hopkins University Press, Baltimore, 2 volumes, 2141 pp.
- Woods, J. L., Harland, D. P., Vernon, J. A., Krsinic, G. L. & Walls, R. J. 2011. Morphology and ultrastructure of Antler velvet hair and body hair from Red Deer (*Cervus elaphus*). *Journal of Morphology* **272**: 34–49. <https://doi.org/10.1002/jmor.10893>
- Xu, X., Zhou, Z. H. & Prum, R. O. 2001. Branched integumental structures in *Sinornithosaurus* and the origin of feathers. *Nature* **410**(6825): 200–204. <https://doi.org/10.1038/35065589>
- Yates, B. C., Espinoza, E. O. & Baker, B. W. 2010. Forensic species identification of elephant (Elephantidae) and giraffe (Giraffidae) tail hair using light microscopy. *Forensic Science, Medicine and Pathology* **6**: 165–171. <https://doi.org/10.1007/s12024-010-9169-6>
- Zazula, G. D., MacKay, G., Andrews, Th., Shapiro, B., Letts, B. & Brock, F. 2009. A late Pleistocene steppe bison (*Bison priscus*) partial carcass from Tsiigehtchic, Northwest Territories, Canada. *Quaternary Science Reviews* **28**: 2734–2742. <https://doi.org/10.1016/j.quascirev.2009.06.012>
- Zima, J. & Andrea, M. 1996. A synopsis of the mammals of the Czech Republic. *Hystrix* **8**(1–2): 107–111.
- Zink, R. M. & Remsen, Jr. J. V. 1986. Evolutionary processes and patterns of geographic variation in birds. *Current Ornithology* **4**: 1–69.

Links:

www.iucnredlist.org – The IUCN Red List of Threatened Species

<https://alaskafurid.wordpress.com>

<http://archive.org/stream/mobot31753003487128#page/9/mode/2up> – Tome Troisième 1867. (ed. Guérin, L.) Nouvelles Archives du Muséum d'Histoire Naturelle de Paris p. 85–116; p. 302.